Fibreoptic control of the laryngeal tube position (Vollmer *et al.*) 2002 19, 306–307
From which countries do chairpersons and invited speakers at important anaesthesia and intensive care meetings come? (Gelb) 2002 19, 75
General anaesthesia for a child with Rubinstein–Taybi syndrome (Tokarz *et al.*) 2002 19, 896–897
General anaesthesia for emergency surgery during acute organophosphate poisoning (Goonasekera *et al.*) 2002 19, 150–152
General anaesthesia with remifentanil and propofol for a patient with centronuclear (myotubular) myopathy (Tokarz *et al.*) 2002 19, 842–844
Gum elastic bougie, capnography and apnoeic oxygenation (Green) 2002 19, 381–382
Hysteroscopic syndrome (Kayacan *et al.*) 2002 19, 770–772
Insidious hypothyroidism unmasked after operation (Vretzakis *et al.*) 2002 19, 532–534
Intravenous regional blocks with guanethidine and prilocaine combined with physiotherapy: two children with complex regional pain syndrome, Type 1 (di Vadi *et al.*) 2002 19, 384–386
Investigating the usefulness of the laryngeal tube in spontaneously breathing patients (Schultz & Buhre) 2002 19, 303–304
Is traditional fasting policy changing? Results of a Swiss survey on current practice (Hofer *et al.*) 2002 19, 894–896
Localized abdominal pain following sympathetic blockade with bretylium for the management of complex regional pain syndrome (Molyneux *et al.*) 2002 19, 147–148
Midodrine, an alternative to intravenous vasopressor therapy after spinal surgery (O'Donnell & Synnott) 2002 19, 841–842
Minor haemodynamic responses in coronary bypass patients following tracheal intubation using either a lightwand device (Trachlight®) or direct laryngoscopy (Vainionpää *et al.*) 2002 19, 229–231
Monitoring in neuroanaesthesia: update of clinical usefulness (Mélot *et al.*) 2002 19, 381
Morbidity in electroconvulsive therapy (Meignan) 2002 19, 614–615
Morbidity in electroconvulsive therapy (Wild *et al.*) 2002 19, 616
Multicompartment block in a parturient undergoing urgent Caesarean section (Sanders & Wee) 2002 19, 226–228
Oasis or mirage? The safety of outpatient dental anaesthesia in hospital (Wildsmith) 2002 19, 762
Peroperative opening of a patent foramen ovale caused by pulmonary artery obstruction due to knotted pulmonary artery catheter (Tavan *et al.*) 2002 19, 145–146
Postoperative hypoxaemia in a patient with patent foramen ovale (Papadopoulos *et al.*) 2002 19, 152–154
Pulmonary oedema after venous air embolism during craniotomy (El Kettani *et al.*) 2002 19, 846–848
Randomizing patients to permit the development of perioperative hypothermia is inappropriate (Booth *et al.*) 2002 19, 688
Remifentanil and acute intermittent porphyria (Durmus *et al.*) 2002 19, 839–840
Salvage with a haemofiltration technique unsuitable in hip prosthesis surgery due to free haemoglobin (Tylman *et al.*) 2002 19, 766–768
Sciatic nerve block and the improvement of femoral nerve block analgesia after total knee replacement (Weber *et al.*) 2002 19, 834–836
Sigmoid capnometry in abdominal aortic aneurysm surgery (Björck) 2002 19, 760–761
Simple preoperative assessment to reduce the risk of traumatic epistaxis during nasotracheal intubation (Kayarkar *et al.*) 2002 19, 690–691
Spontaneous restoration of decreased systemic vascular resistance after spinal anaesthesia (Kamenik) 2002 19, 848–850
Systemic heparinization during postoperative pulmonary embolism induces fatal complications (Kotani *et al.*) 2002 19, 382–384
Thoracic epidurals and avoidance of potential neurological damage (Tucker *et al.*) 2002 19, 772–775
Tongue piercing in an eclamptic patient (Nowicki & Bull) 2002 19, 844–845
Transient respiratory compromise after infraclavicular vertical brachial plexus blockade (Heid *et al.*) 2002 19, 693–694
Ultrasound-guided placement of intra-aortic balloon pump (Šustic *et al.*) 2002 19, 149–150
Use of hyperbaric oxygen for hepatic artery thrombosis following adult orthotopic liver transplantation (Dubost *et al.*) 2002 19, 223–224
Venous gas embolism during liver surgery with argon-enhanced coagulation (Ousmane *et al.*) 2002 19, 225
Ventricular asystole and complete heart block after thoracic epidural analgesia for thymectomy (Lin *et al.*) 2002 19, 160–162
Workforce and regional distribution of anaesthesiologists in Japan (Taga *et al.*) 2002 19, 530–532

Editorials

Acid–base management – is it relevant for the study design of hypothermic neuroprotection? (Frietsch *et al.*) 2002 19, 389–394
Cardiothoracic anaesthesia and the *European Journal of Anaesthesiology* (Feneck) 2002 19, 465
Cholinesterase estimations revisited: the clinical relevance (Karalliedde) 2002 19, 313–316
End-of-life decisions in The Netherlands (Booij) 2002 19, 161–165
Interaction of phenol derivatives with ion channels (Haeseler & Leuwer) 2002 19, 1–8
Liver surgery has become simpler (Franco) 2002 19, 777–779
Murder and the ICU (Park & Khan) 2002 19, 621–623
Oasis or mirage? The safety of outpatient dental anaesthesia in hospital (Bricker) 2002 19, 85–87
Ropivacaine and bupivacaine in obstetric analgesia (Capogna) 2002 19, 237–239
The Cochrane 1998 Albumin Review – not all it was cracked up to be (Horsey) 2002 19, 701–704
Transcranial cerebral oximetry, transcranial Doppler sonography, and heart rate variability: useful neuromonitoring tools in anaesthesia and intensive care? (Schwarz & Litscher) 2002 19, 543–549

Original articles

Accumulation of *S*(+) enantiomer in human beings after general anaesthesia with isoflurane racemate (Haeberle *et al.*) 2002 19, 641–646
Acute normovolaemic haemodilution beyond a haematocrit of 25%: ratio of skeletal muscle tissue oxygen tension and cardiac index is not maintained in the healthy dog (Freitag *et al.*) 2002 19, 487–494
Airway management by physicians wearing anti-chemical warfare gear: comparison between laryngeal mask airway and endotracheal intubation (Goldik *et al.*) 2002 19, 166–169
Alcohol-associated admissions to an adult intensive care unit: an audit (Mostafa & Murthy) 2002 19, 193–196
Anaesthesia for advanced rectal cancer patients treated with combined major resections and intraoperative radiotherapy (Mannaerts *et al.*) 2002 19, 742–748
Anaesthesia manpower in Belgium until 2020: nurse anaesthetists – a possible solution? (Demeere) 2002 19, 755–759
Anaesthetic and postoperative analgesic effects of spinal clonidine as an additive to prilocaine in the transurethral resection of urinary bladder tumours (Santiveri *et al.*) 2002 19, 589–593
Anaesthetic management and outcome in right-lobe living liver-donor surgery (Cammu *et al.*) 2002 19, 93–98

Application of semilinear canonical correlation to the measurement of the electroencephalographic effects of volatile anaesthetics (Bruhn *et al.*) 2002 19, 735–741
Auditory function following spinal analgesia. Comparison of two spinal needles (Malhotra *et al.*) 2002 19, 69–72
Auditory- and somatosensory-evoked potentials in cerebral malaria and anaesthesia: a comparison (Thornton *et al.*) 2002 19, 717–726
Augmented spontaneous breathing and pulmonary gas exchange during pneumoperitoneum (Loeckinger *et al.*) 2002 19, 424–427
Bacterial contamination of propofol: the effects of temperature and lidocaine (Aydin *et al.*) 2002 19, 455–458
Bilateral paravertebral somatic nerve block for ventral hernia repair (Naja *et al.*) 2002 19, 197–202
Can computers help in assessing children's postoperative pain? Initial validation of a computer-assisted interview (Watson *et al.*) 2002 19, 510–516
Cardiac output assessed by arterial thermodilution during exsanguination and fluid resuscitation: experimental validation against a reference technique (Friedman *et al.*) 2002 19, 337–340
Clinical perception of phantom limb sensation in patients with brachial plexus block (Gentili *et al.*) 2002 19, 105–108
Comparison between nasal and intravenous desmopressin for the treatment of aminosalicylic acid-induced platelet dysfunction (Schulz-Stübner *et al.*) 2002 19, 647–651
Comparison of bupivacaine 0.2% and ropivacaine 0.2% combined with fentanyl for epidural analgesia during labour (Aşik *et al.*) 2002 19, 263–270
Comparison of different priming techniques on the onset time and intubating conditions of rocuronium (Yavaşcaoğlu *et al.*) 2002 19, 517–521
Comparison of effects of remifentanil, alfentanil and fentanyl on cardiovascular responses to tracheal intubation in morbidly obese patients (Salihoglu *et al.*) 2002 19, 125–128
Comparison of the effects of remifentanil or fentanyl on anaesthetic induction characteristics of propofol, thiopental or etomidate (Wilhelm *et al.*) 2002 19, 350–356
Conditions for insertion of the laryngeal mask airway: comparisons between sevoflurane and propofol using fentanyl as a co-induction agent. A pilot study (Ganatra *et al.*) 2002 19, 371–375
Cost-reduction analysis of propofol versus sevoflurane: manitenance of anaesthesia for gynaecological surgery using the bispectral index (Struys *et al.*) 2002 19, 727–734
Cuffed oropharyngeal airway as a suitable alternative to the laryngeal mask airway for minor outpatient surgery (Yavaşcaoğlu *et al.*) 2002 19, 203–207
Decrease in minimum alveolar concentration of sevoflurane during anaesthesia and arthroscopy (Higuchi & Adachi) 2002 19, 600–603
Delivery performance of commercial target-controlled infusion devices with Diprifusor™ module (Schraag & Flaschar) 2002 19, 557–560
Does ibuprofen increase perioperative blood loss during hip arthroplasty? (Slappendel *et al.*) 2002 19, 829–851
Doppler indices of diastolic transmitral flow velocity are invalid indicators of myocardial ischaemia during high-dose dobutamine infusion in anaesthetized patients (Filipovic *et al.*) 2002 19, 789–795
Effect of aminophylline on recovery from sevoflurane anaesthesia (Turan *et al.*) 2002 19, 452–454
Effect of different doses of cisatracurium on intraocular pressure in sedated patients (Sator-Katzenschlager *et al.*) 2002 19, 823–828
Effect of three different doses of urapidil on blood glucose concentrations in the streptozotocin diabetic rat (Ittner *et al.*) 2002 19, 504–509
Effect of VIMA with sevoflurane versus TIVA with propofol or midazolam-sufentanil on the cytokine response during CABG surgery (El Azab *et al.*) 2002 19, 276–282
Effects of adding alfentanil or atracurium to lidocaine solution for intravenous regional anaesthesia (Kurt *et al.*) 2002 19, 522–525
Effects of chronic occupational exposure to anaesthetic gases on the rate of neutrophil apoptosis among anaesthetists (Tyther *et al.*) 2002 19, 604–608
Effects of clonidine and dopamine on gastric tone (Levein *et al.*) 2002 19, 99–104
Effects of induction of anaesthesia with sufentanil and positive-pressure ventilation on the intra- to extrathoracic volume distribution (von Spiegel *et al.*) 2002 19, 428–435
Effects of normovolaemic haemodilution on middle cerebral artery blood flow velocity and oxygen delivery (Karadibak *et al.*) 2002 19, 330–336
Effects of sodium nitroprusside-induced controlled hypotension on pancreatic function assessed by pancreatitis-associated protein in patients undergoing radical prostatectomy (Piper *et al.*) 2002 19, 609–613
Effects of volume and concentration of lidocaine on epidural anaesthesia in pregnant females (Nakayama *et al.*) 2002 19, 808–811
Epidural bupivacaine with sufentanil or fentanyl during labour: a randomized, double-blind study (Rolfseng *et al.*) 2002 19, 812–818
Epidural combination of ropivacaine with sufentanil for postoperative analgesia after total knee replacement: a pilot study (Kampe *et al.*) 2002 19, 666–671
Epidural morphine injection after combined spinal and epidural anaesthesia (Takenaka-Hamaya *et al.*) 2002 19, 672–676
Fate of abstracts from the Paris 1995 European Society of Anaesthesiologists meeting (Castillo *et al.*) 2002 19, 888–895
Fentanyl added to bupivacaine 0.05% or ropivacaine 0.05% in patient-controlled epidural analgesia in labour (Pirbudak *et al.*) 2002 19, 271–275
First clinical experience with the rapid-, short-acting amiodarone derivative E 047/1 after cardiac surgery (Gombotz *et al.*) 2002 19, 23–31
Fluoride ion toxicity in rabbit kidney thick ascending limb cells (Cittanova *et al.*) 2002 19, 341–349
Flurbiprofen does not change the bispectral index and 95% spectral edge frequency during total intravenous anaesthesia with propofol and fentanyl (Hirota *et al.*) 2002 19, 483–486
Gastric emptying in post-thoracotomy patients receiving a thoracic fentanyl-bupivacaine epidural infusion (Guha *et al.*) 2002 19, 652–657
Haemodynamic and splanchnic organ blood flow responses during sevoflurane-induced hypotension in dogs (Takeda *et al.*) 2002 19, 442–446
Haemodynamic and splanchnic organ blood-flow responses to hypotension induced by the calcitonin gene-related peptide and sodium nitroprusside in dogs (Takeda & Tomaru) 2002 19, 876–882
Haemodynamic changes in ischaemic vs. anhepatic pig experimental model of acute liver failure (Theodoraki *et al.*) 2002 19, 40–46
Impact of the reduction of anaesthesia turnover time on operating room efficiency (Sokolovic *et al.*) 2002 19, 560–563
In vitro effects of 4-chloro-3-ethylphenol in skeletal muscle preparations from malignant hyperthermia susceptible and normal swine (Gerbershagen *et al.*) 2002 19, 135–140
Inhalation anaesthesia is cost-effective for ambulatory surgery: a clinical comparison with propofol during elective knee arthroscopy (Dolk *et al.*) 2002 19, 88–92
Intraoperative fetal oxygen saturation during Caesarean section: general anaesthesia using sevoflurane with either 100% oxygen or 50% nitrous oxide in oxygen (Parpaglioni *et al.*) 2002 19, 115–118
Intravenous single-dose tramadol versus meperidine for pain relief in renal colic (Eray *et al.*) 2002 19, 368–370

Iontophoretically applied lidocaine and the prevention of pain associated with the injection of intravenous propofol: a comparison with intravenous lidocaine (Sharpe *et al.*) 2002 19, 170–172
Ketamine and the inhibition of albumin extravasation in chemical peritonitis in rat (Hirota *et al.*) 2002 19, 173–176
Ketamine effects on human neuronal Na^+ channels (Reckziegel *et al.*) 2002 19, 634–640
Lidocaine-induced cell death in a human model of neuronal apoptosis (Friederich & Schmitz) 2002 19, 564–570
Magnesium infusion reduces perioperative pain (Kara *et al.*) 2002 19, 52–56
Manipulations in glycogen metabolism and the failure to influence infarct size in the ischaemic rabbit heart (Gozal *et al.*) 2002 19, 495–503
Monitoring intravascular volumes for postoperative volume therapy (Brock *et al.*) 2002 19, 288–294
Myocardial ischaemia in patients with impaired left ventricular function undergoing coronary artery bypass grafting – milrinone versus nifedipine (Möllhoff *et al.*) 2002 19, 796–802
Neuromuscular effects of rapacuronium on the diaphragm and skeletal muscles in anaesthetized patients using cervical magnetic stimulation for stimulating the phrenic nerves (Moerer *et al.*) 2002 19, 883–887
Operating room management and strategies in Switzerland: results of a survey (Sieber & Leibundgut) 2002 19, 415–423
Oxygen transport and myocardial function after the administration of albumin 5%, hydroxyethylstarch 6% and succinylated gelatine 4% to rabbits (Himpe *et al.*) 2002 19, 860–867
Perioperative risk factors in elective pneumonectomy: the impact of excess fluid balance (Møller *et al.*) 2002 19, 57–62
Posterior subgluteal approach to block the sciatic nerve: description of the technique and initial clinical experiences (Di Benedetto *et al.*) 2002 19, 682–686
Postoperative pain management in orthopaedic patients: no differences in pain score, but improved stress control by epidural anaesthesia (Adams *et al.*) 2002 19, 658–665
Postoperative residual curatization with cisatracurium and rocuronium infusions (Cammu *et al.*) 2002 19, 129–134
Predictors of successful neuraxial block: a prospective study (de Oliveira Filho *et al.*) 2002 19, 447–451
Preload and haemodynamic assessment during liver transplantation: a comparison between the pulmonary artery catheter and transpulmonary indicator dilution techniques (Della Rocca *et al.*) 2002 19, 868–875
Prophylactic intravenous bolus ephedrine for elective Caesarean section under spinal anaesthesia (Loughrey *et al.*) 2002 19, 63–68
Propofol in low doses causes redistribution of body heat in male volunteers (Noguchi *et al.*) 2002 19, 677–681
Propofol in rats: testing for nonlinear pharmacokinetics and modelling acute tolerance to EEG effects (Ihmsen *et al.*) 2002 19, 177–188
Psychomotor performance after short-term anaesthesia (Haavisto & Kauranen) 2002 19, 436–441
QT interval and QT dispersion during the induction of anaesthesia in patients with subarachnoid haemorrhage: a comparison of thiopental and propofol (Tanskanen *et al.*) 2002 19, 749–754
Randomized comparison of remifentanil–propofol with a sciatic–femoral nerve block for out-patient knee arthroscopy (Casati *et al.*) 2002 19, 109–114
Randomized controlled trial investigating the effect of transcervical papaverine and bupivacaine on postoperative analgesia following laparoscopic sterilization (Ng *et al.*) 2002 19, 803–807
Sedation assessment in critically ill patients with bispectral index (Riess *et al.*) 2002 19, 18–22
Shorter discharge time after regional or intravenous anaesthesia in combination with laryngeal mask airway compared with balanced anaesthesia with endotracheal intubation (Junger *et al.*) 2002 19, 119–124
Spinal block or total intravenous anaesthesia with propofol and remifentanil for gynaecological outpatient procedures (Danelli *et al.*) 2002 19, 594–599
Stability of a sufentanil–ropivacaine mixture in a glass and a PVC reservoir (Brodner *et al.*) 2002 19, 295–297
Subcutaneous epinephrine administration decreases lower oesophageal sphincter pressure and gastro-oesophageal pressure gradient in children under general anaesthesia (Kohjitani & Obara) 2002 19, 189–192
Succinylcholine improves the time of performance of a tracheal intubation in prehospital critical care medicine (Ricard-Hibon *et al.*) 2002 19, 361–367
Sufentanil or clonidine for blunting the increase in intraocular pressure during rapid-sequence induction (Georgiou *et al.*) 2002 19, 819–822
Target-controlled propofol requirements at induction of anaesthesia: effect of remifentanil and midazolam (Conway *et al.*) 2002 19, 580–584
The effects of fenoldopam on renal function in patients undergoing elective aortic surgery (Halpenny *et al.*) 2002 19, 32–39
The prevention of propofol injection pain by tramadol or ondansetron (Memiş *et al.*) 2002 19, 47–51
Thiopental impairs neutrophil oxidative response by inhibition of intracellular signalling (Fröhlich *et al.*) 2002 19, 474–482
Use of sevoflurane during cardiopulmonary bypass decreases incidence of awareness (Celebioğlu *et al.*) 2002 19, 283–287
Using the Sustained Attention to Response Test to detect cognitive dysfunction after day case surgery (Thompson *et al.*) 2002 19, 585–588
Voltage-dependent block of neuronal and skeletal muscle sodium channels by thymol and menthol (Haeseler *et al.*) 2002 19, 571–579

Reviews

Allergic reactions occurring during anaesthesia (Mertes & Laxenaire) 2002 19, 240–262
Anaesthesia for elective liver resection: some points should be revisited (Lentschener & Ozier) 2002 19, 780–788
Anaesthetic agents in paediatric day case surgery: do they affect outcome? (Moore *et al.*) 2002 19, 9–17
Cardiac arrest, chain of survival and Utstein style (Gullo) 2002 19, 624–633
Central neuraxial blocks and anticoagulation: a review of current trends (Tyagi & Bhattacharya) 2002 19, 317–329
Isoprostanes – markers of ischaemia reperfusion injury (Sakamoto *et al.*) 2002 19, 550–559
Laudanosine, an atracurium and cisatracurium metabolite (Fodale & Santamaria) 2002 19, 466–473
Long QT syndrome and anaesthesia (Wisely & Shipton) 2002 19, 855–859
Neurological complications following central neuraxial blocks: are there predictive factors? (De Tommaso *et al.*) 2002 19, 705–716
Pathophysiology of cocaine abuse (Brownlow & Pappachan) 2002 19, 395–414

European Journal of Anaesthesiology 2002; 19: 908–916
© 2002 European Academy of Anaesthesiology
ISSN 0265-0215

Author Index

Acar, Ş. H. V. *see* Yavaşcaoğlu, B.
Adachi, Y. *see* Higuchi, H.
Adams, H. A., Saatweber, P., Schmitz, C. S. & Hecker, H. Postoperative pain management in orthopaedic patients: no differences in pain score, but improved stress control by epidural anaesthesia, 658
Addison, R. *see* Cunnington, P. M. D.
Adriaensen, H. F. *see* Himpe, D. G.
Agritmis, A. *see* Karaca, S.
Akça, S. *see* Eray, O.
Akcali, D. T. *see* Mahli, A.
Akgün, I. *see* Karaca, S.
Albertin, A. *see* Danelli, G.
Alberts, P. A. *see* Gerbershagen, M. U.
Alkış, N. *see* Aşık, İ.
Alon, E. *see* Hofer, C. K.
Amemiya, Y. *see* Noguchi, I.
Anderson, R. E. *see* Dolk, A.
Annual Meeting of the European Academy of Anaesthesiology 2001, 28 August–1 September 2001: Graz, Austria, 208
Argiriadou, H. *see* Georgiou, M.
Arkadopoulos, N. *see* Theodoraki, K.
Arkan, A. *see* Karadibak, K.
Arkuszewski, P. *see* Tokarz, A.
Arnaoutoglou, E. *see* Papadopoulos, G.
Arnestad, J.-P. *see* Tylman, M.
Arxer, A. *see* Santiveri, X.
Asai, T., Hidaka, I., Kubota, T. & Kawachi, S. Efficacy of the laryngeal tube, 505
Asif, M. *see* Sharpe, P.
Aşık, İ., Göktuğ, A., Gülay, I., Alkış, N. & Uysalel, A. Comparison of bupivacaine 0.2% and ropivacaine 0.2% combined with fentanyl for epidural analgesia during labour, 263
Atar, D. *see* Filipovic, M.
Atcheson, R. A. *see* Thompson, J. P.
Attapattu, C. L. K. *see* Goonasekera, C. D. A.
Auweiler, M. *see* Kampe, S.
Awan, M. *see* O'Brien, B.
Aydin, N. *see* Aydin, O. N.
Aydin, O. N., Aydin, N., Gultekin, B., Ozgun, S. & Gurel, A. Bacterial contamination of propofol: the effects of temperature and lidocaine, 455
Aydoğdu, T. *see* Kara, H.
Aygünes, B. *see* Kurt, N.
Aypar, Ü. *see* Celebioğlu, B.

Baba, H. *see* Taga, K.
Baba, S. *see* Hirota, K.
Badaoui, R. *see* El Kettani, C.
Baller, C. *see* Moerer, O.
Baltaci, B. *see* Kaymak, C.
Bartels, M. *see* Wild, B.
Ben-Abraham, R. *see* Goldik, Z.
Bengtson, J.-P. *see* Tylman, M.
Bengtsson, A. *see* Tylman, M.
Benraad, B. *see* Slappendel, R.
Benson, M. *see* Junger, A.
Berendes, E. *see* Möllhoff, T.
Berkenstadt, H. *see* Friedman, Z.
Berré, J. *see* Mélot, C.
Berti, M. *see* Casati, A.
Berti, M. *see* Danelli, G.
Bertini, L. *see* Di Benedetto, P.
Bhattacharya, A. *see* Tyagi, A.
Bibl, D. *see* Brock, H.
Biedler, A. *see* Wilhelm, W.
Biro, P. *see* Sokolovic, E.
Björck, M. Sigmoid capnometry in abdominal aortic aneurysm surgery, 760
Blanc, O. *see* Cittanova, M. L.
Blondet, R. *see* Tavan, R.
Boldt, J. *see* Piper, S. N.
Bone, H. G. *see* Riess, M. L.
Bonnet, F. *see* Gentili, M. E.
Booij, L. H. D. J. End-of-life decisions in The Netherlands, 161
Booth, M. G., Gallagher, G. & Kinsella, J. Randomizing patients to permit the development of perioperative hypothermia is inappropriate, 688
Borchgrevink, P. C. *see* Rolfseng, O. K.
Bornstein, J. *see* Goldik, Z.
Boucher-Hayes, D. *see* Halpenny, M.
Bouillon, T. *see* Bruhn, J.
Bourbouze, R. *see* Cittanova, M. L.
Brambrink, A. M. *see* Heid, F. M.
Breen, P. *see* Halpenny, M.
Bricker, S. Oasis or mirage? The safety of outpatient dental anaesthesia in hospital, 85
Bricker, S. A reply, 763
Brill, S. *see* di Vadi, P. P.
Brock, H., Gabriel, C., Bibl, D. & Necek, S. Monitoring intravascular volumes for postoperative volume therapy, 288
Brodner, G., Ermert, T., Van Aken, H., Westphal, M., Marcus, M. A. E., Gogarten, W., Goeters, C. & Bürkle, H. Stability of a sufentanil–ropivacaine mixture in a glass and a PVC reservoir, 295
Brown, C. *see* di Vadi, P. P.
Brownlow, H., Sutherland, P. & Lanham, P. An unusual case of breathing circuit obstruction, 155
Brownlow, H. A. & Pappachan, J. Pathophysiology of cocaine abuse, 395
Bruhn, J., Rehberg, B., Röpcke, H., Bouillon, T. & Hoeft, A. Application of semilinear canonical correlation to the measurement of the electroencephalographic effects of volatile anaesthetics, 735
Bücheler, O. *see* Wilhelm, W.
Bucher, M. *see* Ittner, K. P.
Buerkle, H. *see* Möllhoff, T.
Bufler, J. *see* Haeseler, G.
Bugter, M. L. T. *see* Slappendel, R.
Buhre, W. *see* Schultz, U.
Bull, P. T. *see* Nowicki, R. W. A.
Bürkle, H. *see* Brodner, G.
Buscher, H. *see* Moerer, O.
Buser, P. *see* Filipovic, M.
Butani, M. *see* D'Mello, J.
Butani, M. *see* Ganatra, S. B.

Calam, R. *see* Watson, S.
Cammu, G., de Baerdemaeker, L., den Blauwen, N., de Mey, J.-C., Struys, M. & Mortier, E. Postoperative residual curarization with cisatracurium and rocuronium infusions, 129
Cammu, G., Troisi, R., Cuomo, O., de Hemptinne, B., Di Florio, E. & Mortier, E. Anaesthetic management and outcome in right-lobe living liver-donor surgery, 93
Cannerfelt, R. *see* Dolk, A.
Capogna, G. *see* Parpaglioni, R.
Capogna, G. Ropivacaine and bupivacaine in obstetric analgesia, 257
Caponas, G. & Popat, M. Difficulty in use and device failure with the intubating laryngeal mask airway®, 764
Caporuscio, A. *see* De Tommaso, O.
Cappelleri, G. *see* Casati, A.
Carrero-Cardenal, E. J. Differentiating massive carbon dioxide embolism from tension pneumothorax during laparoscopic cholecystectomy, 459
Casati, A. *see* Danelli, G.
Casati, A. *see* Di Benedetto, P.
Casati, A., Cappelleri, G., Berti, M., Fanelli, G., Di Benedetto, P. & Torri, G. Randomized comparison of remifentanil–propofol with a sciatic–femoral nerve block for out-patient knee arthroscopy, 109
Castillo, J. Garcia-Guasch, R. & Cifuentes, I. Fate of abstracts from the Paris 1995 European Society of Anaesthesiologists meeting, 888
Cebelli, V. *see* Yavaşcaoğlu, B.
Celebioğlu, B., Pamuk, A. G., Aypar, Ü. & Paşaoğlu, I. Use of sevoflurane during cardiopulmonary bypass decreases incidence of awareness, 283
Celebioğlu, B. & Yener, F. Anaesthesia for open-heart surgery in a patient with Weaver's syndrome, 897
Çelik, Ç. *see* Pirbudak, L.
Celleno, D. *see* Parpaglioni, R.
Çete, N. *see* Eray, O.
Çete, Y. *see* Eray, O.
Cherng, C.-H. *see* Lin, T.-C.
Chevanne, F. *see* Estebe, J. P.
Chiefari, M. *see* Iannuzzi, E.
Chollet, C. *see* Ricard-Hibon, A.
Cifuentes, I. *see* Castillo, J.
Cirillo, V. *see* Iannuzzi, E.
Cittanova, M. L., Estepa, L., Bourbouze, R., Blanc, O., Verpont, M.-C., Wahbe, F., Coriat, P., Daudon, M. & Ronco, P. M. Fluoride ion toxicity in rabbit kidney thick ascending limb cells, 341
Coccia, C. *see* Della Rocca, G.
Çolak, A. *see* Turan, A.
Conway, D. H., Hasan, S. K. & Simpson, M. E. Target-controlled propofol requirements at induction of anaesthesia: effect of remifentanil and midazolam, 580
Corcoran, T. B. *see* Sakamoto, H.
Coriat, P. *see* Cittanova, M. L.
Coronel, B. *see* Tavan, R.
Coskun, D. *see* Mahli, A.
Costa, M. G. *see* Della Rocca, G.
Cranfield, K. *see* Kampe, S.
Crozier, T. A. *see* Moerer, O.
Cunningham, A. J. *see* Halpenny, M.
Cunnington, P. M. D. & Addison, R. Dilated cardiomyopathy in dystrophic epidermolysis bullosa: a lethal complication of epidermolysis bullosa, 689
Cuomo, O. *see* Cammu, G.
Curtis, N. *see* Thornton, C.

D'Mello, J., Pagedar, R., Butani, M., Kurkal, P. & Pandey, K. Use of the intubating laryngeal mask airway in a case of ankylosing spondylitis for coronary artery bypass grafting, 298
D'Mello, J. *see* Ganatra, S. B.
da Fonseca, M. H. Z. *see* de Oliveira Filho, G. R.
Danelli, G., Berti, M., Casati, A., Albertin, A., Deni, E., Nobili, F. & Torri, G. Spinal block or total intravenous anaesthesia with propofol and remifentanil for gynaecological outpatient procedures, 594
Daudon, M. *see* Cittanova, M. L.
Davidson, A. C. *see* Ng, A.
de Baerdemaeker, L. *see* Cammu, G.
de Hemptinne, B. *see* Cammu, G.
De Hert, S. G. *see* Himpe, D. G.
De Lange, J. J. *see* El Azab, S. R.
de Mey, J.-C. *see* Cammu, G.
de Oliveira Filho, G. R., Gomes, H. P., da Fonseca, M. H. Z., Hoffman, J. C., Pederneiras, S. G. & Garcia, J. H. S. Predictors of successful neuraxial block: a prospective study, 147
De Tommaso, O., Caporuscio, A. & Tagariello, V. Neurological complications following central neuraxial blocks: are there predictive factors?, 705
Della Rocca, G., Costa, M. G., Coccia, C., Pompei, L. & Pietropaoli, P. Preload and haemodynamic assessment during liver transplantation: a comparison between the pulmonary artery catheter and transpulmonary indicator dilution techniques, 868
Demeere, J. L. Anaesthesia manpower in Belgium until 2020: nurse anaesthetists – a possible solution?, 755
Demirbilek, S. *see* Togal, T.
Demirkiran, O. *see* Salihoglu, Z.
Demiroluk, S. *see* Salihoglu, Z.
den Blauwen, N. *see* Cammu, G.
Deng, M. C. *see* Möllhoff, T.
Dengler, R. *see* Haeseler, G.
Deni, E. *see* Danelli, G.
Dergeloo, O. *see* Filipovic, M.
Di Benedetto, P. *see* Casati, A.
Di Benedetto, P., Casati, A., Bertini, L. & Fanelli, G. Posterior subgluteal approach to block the sciatic nerve: description of the technique and initial clinical experiences, 682
Di Florio, E. *see* Cammu, G.
di Vadi, P. P., Brill, S., Jack, T., Brown, C. & Edwards, T. Intravenous regional blocks with guanethidine and prilocaine combined with physiotherapy: two children with complex regional pain syndrome, Type 1, 384
Dietenbach, C. *see* Kampe, S.
Dieterich, H.-J. *see* Haeberle, H. A.
Dirksen, R. *see* Slappendel, R.
Dohi, S. *see* Takenaka-Hamaya, C.
Dolk, A., Cannerfelt, R., Anderson, R. E. & Jakobsson, J. Inhalation anaesthesia is cost-effective for ambulatory surgery: a clinical comparison with propofol during elective knee arthroscopy, 88
Dollo, G. *see* Estebe, J. P.
Döşemeci, L. *see* Kayacan, N.
Dubost, T., Goubaux, B., Duhalde, M., Raucoules-Aime, M., Wolkiewiez, J. & Gugenheim, J. Use of hyperbaric oxygen for hepatic artery thrombosis following adult orthotopic liver transplantation, 223
Duhalde, M. *see* Dubost, T.
Duperret, S. *see* Tavan, R.
Durmus, M. *see* Togal, T.
Durmus, M., Turkoz, A., Togal, T., Koroglu, A., Toprak, H. I. & Ersoy, M. O. Remifentanil and acute intermittent porphyria, 839
Dutta, A. *see* Malhotra, S. K.
Dyzers, D. *see* Struys, M. M. R. F.

Ecoffey, C. *see* Estebe, J. P.
Eden, A. *see* Goldik, Z.

Edwards, T. *see* di Vadi, P. P.
El Azab, S. R., Rosseel, P. M. J., De Lange, J. J., van Wijk, E. M., van Strik, R. & Scheffer, G. J. Effect of VIMA with sevoflurane versus TIVA with propofol or midazolam-sufentanil on the cytokine response during CABG surgery, 276
El Kettani, C., Badaoui, R., Fikri, M., Jeanjean, P., Montpellier, D. & Tchaoussoff, J. Pulmonary oedema after venous air embolism during craniotomy, 846
Ellinger, K. *see* Vollmer, T.
Elliott, R. E. *see* Moore, E. W.
Engquist, A. *see* Møller, A. M.
Eray, O., Çete, Y., Oktay, C., Karslı, B., Akça, S., Çete, N. & Ersoy, F. Intravenous single-dose tramadol versus meperidine for pain relief in renal colic, 368
Erbayraktar, S. *see* Karadibak, K.
Erkan, N. *see* Karadibak, K.
Erman, M. *see* Kayacan, N.
Ermert, T. *see* Brodner, G.
Errando, C. L. Anaesthesia for the professional singer, 687
Ersoy, F. *see* Eray, O.
Ersoy, M. O. *see* Durmus, M.
Ersoy, O. *see* Togal, T.
Eschweiler, G. W. *see* Wild, B.
Estebe, J. P., Ecoffey, C., Dollo, G., Le Corre, P., Chevanne, F. & Le Verge, R. Bupivacaine pharmacokinetics and motor blockade following epidural administration of the bupivacaine-sulphobutylether 7-β-cyclodextrin complex in sheep, 308
Estepa, L. *see* Cittanova, M. L.

Fanelli, G. *see* Casati, A.
Fanelli, G. *see* Di Benedetto, P.
Feneck, R. Cardiothoracic anaesthesia and the *European Journal of Anaesthesiology*, 465
Ferdi, E. *see* Vretzakis, G.
Fiege, M. *see* Gerbershagen, M. U.
Fikri, M. *see* El Kettani, C.
Filipovic, M., Seeberger, M. D., Rohlfs, R., Dergeloo, O., Studer, W., Atar, D., Buser, P. & Skarvan, K. Doppler indices of diastolic transmitral flow velocity are invalid indicators of myocardial ischaemia during high-dose dobutamine infusion in anaesthetized patients, 789
Flaschar, J. *see* Schraag, S.
Fleyfel, M. *see* Ousmane, M. L.
Fodale, V. & Santamaria, L. B. Laudanosine, an atracurium and cisatracurium metabolite, 466
Fournier, R. *see* Weber, A.
Fox, A. J. *see* Thompson, J. P.
Francis, G. A. *see* Kayarkar, R.
Franco, D. Liver surgery has become simpler, 777
Freitag, M., Standl, T., Horn, E.-P., Wilhelm, S. & Schulte am Esch, J. Acute normovolaemic haemodilution beyond a haematocrit of 25%: ratio of skeletal muscle tissue oxygen tension and cardiac index is not maintained in the healthy dog, 487
Friederich, P. *see* Reckziegel, G.
Friederich, P. & Schmitz, T. P. Lidocaine-induced cell death in a human model of neuronal apoptosis, 564
Friedman, Z., Berkenstadt, H., Margalit, N., Segal, E. & Perel, A. Cardiac output assessed by arterial thermodilution during exsanguination and fluid resuscitation: experimental validation against a reference technique, 337
Frietsch, T., Piepgras, A., Krafft, P., Schwab, S., Kuschinsky, W. & Waschke, K. F. Acid–base management – is it relevant for the study design of hypothermic neuroprotection?, 589
Fröhlich, D., Wittmann, S., Rothe, G., Schmitz, G. & Taeger, K. Thiopental impairs neutrophil oxidative response by inhibition of intracellular signaling, 474
Fujihara, H. *see* Taga, K.
Fukui, K., Morioka, T., Hisada, K., Nishio, S., Irita, K. & Takahashi, S. Effects of thiamylal, sevoflurane and isoflurane on the cortically recorded somatosensory evoked potentials, 899
Fukushi, S. *see* Hirota, K.
Fusco, P. *see* Parpaglioni, R.

Gabriel, C. *see* Brock, H.
Gallagher, G. *see* Booth, M. G.
Gamulin, Z. *see* Weber, A.
Ganatra, S. B., D'Mello, J., Butani, M. & Jhamnani, P. Conditions for insertion of the laryngeal mask airway: comparisons between sevoflurane and propofol using fentanyl as a co-induction agent. A pilot study, 571
Garcia, J. H. S. *see* de Oliveira Filho, G. R.
Garcia-Guasch, R. *see* Castillo, J.
Gardiner, J. *see* Loughrey, J. P. R.
Gaszyński, T. *see* Tokarz, A.
Gaszyński, W. *see* Tokarz, A.
Gelb, A. W. From which countries do chairpersons and invited speakers at important anaesthesia and intensive care meetings come?, 75
Gentili, M. E., Verton, C., Kinirons, B. & Bonnet, F. Clinical perception of phantom limb sensation in patients with brachial plexus block, 105
Genzwuerker, H. V. *see* Vollmer, T.
Georgiou, M., Parlapani, A., Argiriadou, H., Papagiannopoulou, P., Katsikis, G. & Kaprini, E. Sufentanil or clonidine for blunting the increase in intraocular pressure during rapid-sequence induction, 819
Gerbershagen, M. U., Wappler, F., Fiege, M., Weißhorn, R., Alberts, P. A., von Breunig, F. & Schulte am Esch, J. *In vitro* effects of 4-chloro-3-ethylphenol in skeletal muscle preparations from malignant hyperthermia susceptible and normal swine, 135
Giannaris, S. *see* von Spiegel, T.
Goeters, C. *see* Brodner, G.
Goeters, C. *see* Riess, M. L.
Gogarten, W. *see* Brodner, G.
Gökmen, N. *see* Karadibak, K.
Göksu, S. *see* Pirbudak, L.
Göktay, Y. *see* Karadibak, K.
Göktuğ, A. *see* Aşık, İ.
Goldik, Z., Bornstein, J., Eden, A. & Ben-Abraham, R. Airway management by physicians wearing anti-chemical warfare gear: comparison between laryngeal mask airway and endotracheal intubation, 166
Gombotz, H., Vicenzi, M., Mahla, E., Rehak, P. & Metzler, H. First clinical experience with the rapid-, short-acting amiodarone derivative E 047/1 after cardiac surgery, 23
Gomes, H. P. *see* de Oliveira Filho, G. R.
Goonasekera, C. D. A., Perhiyagoda, C. J. B., Attapattu, C. L. K. & Nagarathne, M. D. General anaesthesia for emergency surgery during acute organophosphate poisoning, 150
Goubaux, B. *see* Dubost, T.
Gozal, Y., Wolff, R. A. & Van Winkle, D. M. Manipulations in glycogen metabolism and the failure to influence infarct size in the ischaemic rabbit heart, 495
Graefe, U. A. *see* Riess, M. L.
Green, D. W. Gum elastic bougie, capnography and apnoeic oxygenation, 381
Grobecker, H. F. *see* Ittner, K. P.
Grosskreutz, J. *see* Haeseler, G.
Grover, S. *see* Malhotra, S. K.
Gugenheim, J. *see* Dubost, T.
Guha, A., Scawn, N. D. A., Rogers, S. A., Pennefather, S. H. & Russell, G. N. Gastric emptying in post-thoracotomy patients receiving a thoracic fentanyl-bupivacaine epidural infusion, 652

Gülay, I. *see* Aşık, İ.
Gulhan, Y. *see* Kaymak, C.
Gullo, A. Cardiac arrest, chain of survival and Utstein style, 624
Gultekin, B. *see* Aydin, O. N.
Gurel, A. *see* Aydin, O. N.

Haavisto, E. & Kauranen, K. Psychomotor performance after short-term anaesthesia, 436
Haavisto, E. *see* Vainionpää, V.
Habib, A. *see* Ng, A.
Haeberle, H. A., Wahl, H. G., Jakubetz, H., Krause, H., Schmidt, R., Schurig, V. & Dieterich, H.-J. Accumulation of *S*(+) enantiomer in human beings after general anaesthesia with isoflurane racemate, 641
Haeseler, G. & Leuwer, M. Interaction of phenol derivatives with ion channels, 1
Haeseler, G., Maue, D., Grosskreutz, J., Bufler, J., Nentwig, B., Piepenbrock, S., Dengler, R. & Leuwer, M. Voltage-dependent block of neuronal and skeletal muscle sodium channels by thymol and menthol, 571
Haisch, G. *see* Piper, S. N.
Haller, U. *see* Sokolovic, E.
Halligan, M. *see* Tyther, R.
Halpenny, M., Rushe, C., Breen, P., Cunningham, A. J., Boucher-Hayes, D. & Shorten, G. D. The effects of fenoldopam on renal function in patients undergoing elective aortic surgery, 32
Hamaya, Y. *see* Takenaka-Hamaya, C.
Hardee, P. S. G. F. *see* Klein, A. A.
Hartmann, B. *see* Junger, A.
Hasan, S. K. *see* Conway, D. H.
Hecker, H. *see* Adams, H. A.
Heid, F. M., Kern, T. & Brambrink, A. M. Transient respiratory compromise after infraclavicular vertical brachial plexus blockade, 693
Heinze, G. *see* Sator-Katzenschlager, S. M.
Hempelmann, G. *see* Junger, A.
Herregods, L. L. G. *see* Struys, M. M. R. F.
Heyderman, R. S. *see* Thornton, C.
Hidaka, I. *see* Asai, T.
Higuchi, H. & Adachi, Y. Decrease in minimum alveolar concentration of sevoflurane during anaesthesia and arthroscopy, 600
Himpe, D. G., De Hert, S. G., Vermeyen, K. M. & Adriaensen, H. F. Oxygen transport and myocardial function after the administration of albumin 5%, hydroxyethylstarch 6% and succinylated gelatine 4% to rabbits, 860
Hinz, J. *see* Moerer, O.
Hirota, K., Fukushi, S., Baba, S. & Matsuki, A. Flurbiprofen does not change the bispectral index and 95% spectral edge frequency during total intravenous anaesthesia with propofol and fentanyl, 483
Hirota, K., Ishihara, H. & Matsuki, A. Ketamine and the inhibition of albumin extravasation in chemical peritonitis in rat, 173
Hisada, K. *see* Fukui, K.
Hoeft, A. *see* Bruhn, J.
Hoeft, A. *see* von Spiegel, T.
Hoermann, C. *see* Loeckinger, A.
Hofer, C. K., Zollinger, A., Tucci, M., van Gessel, E., Alon, E. & Klaghofer, R. Is traditional fasting policy changing? Results of a Swiss survey on current practice, 894
Hoffman, J. C. *see* de Oliveira Filho, G. R.
Horn, E.-P. *see* Freitag, M.
Horsey, P. J. The Cochrane 1998 Albumin Review – not all it was cracked up to be, 701
Hsu, C.-H. *see* Lin, T.-C.
Huppert, A. *see* Wilhelm, W.

Iannuzzi, E., Iannuzzi, M., Pedicini, M. S., Cirillo, V., Chiefari, M. & Sacerdoti, G. Anaphylactic reaction after cisatracurium administration, 691
Iannuzzi, M. *see* Iannuzzi, E.
Ichinose, H. *see* Nakayama, M.
Ihmsen, H., Tzabazis, A., Schywalsky, M. & Schwilden, H. Propofol in rats: testing for nonlinear pharmacokinetics and modelling acute tolerance to EEG effects, 177
Irita, K. *see* Fukui, K.
Ishihara, H. *see* Hirota, K.
Ishihara, H. *see* Kotani, N.
Ishihara, H. *see* Tsubo, T.
Ittner, K. P., Bucher, M., Zimmermann, M., Grobecker, H. F., Krämer, B. K. & Taeger, K. Effect of three different doses of urapidil on blood glucose concentrations in the streptozotocin diabetic rat, 504

Jack, T. *see* di Vadi, P. P.
Jakobsson, J. *see* Dolk, A.
Jakubetz, H. *see* Haeberle, H. A.
Jankovic, Z., Radojevic, D., Neskovic, V. & Stamenkovic, D. Acute abdomen after coronary artery bypass surgery masked by thoracic epidural analgesia, 526
Jeanjean, P. *see* El Kettani, C.
Jhamnani, P. *see* Ganatra, S. B.
Jimmieson, P. *see* Watson, S.
Joshi, M. *see* Malhotra, S. K.
Junger, A., Klasen, J., Hartmann, B., Benson, M., Röhrig, R., Kuhn, D. & Hempelmann, G. Shorter discharge time after regional or intravenous anaesthesia in combination with laryngeal mask airway compared with balanced anaesthesia with endotracheal intubation, 119

Kahveci, Ş. F. *see* Yavaşçaoğlu, B.
Kamenik, M. Spontaneous restoration of decreased systemic vascular resistance after spinal anaesthesia, 848
Kampe, S., Diefenbach, C., Kanis, B., Auweiler, M., Kiencke, P. & Cranfield, K. Epidural combination of ropivacaine with sufentanil for postoperative analgesia after total knee replacement: a pilot study, 666
Kanaya, N. *see* Nakayama, M.
Kanis, B. *see* Kampe, S.
Kaprini, E. *see* Georgiou, M.
Kara, H., Şahin, N., Ulusan, V. & Aydoğdu, T. Magnesium infusion reduces perioperative pain, 52
Karaaslan, K. *see* Togal, T.
Karaca, S. & Akgün, I. Epidural anaesthesia for arthroscopic knee surgery in a patient with Takayasu's arteritis, 528
Karaca, S., Akgün, I. & Agritmis, A. Epidural anaesthesia in a patient with POEMS syndrome, 154
Karaca, S. & Ünlüsoy, E. Ö. Accidental injection of intravenous bupivacaine, 616
Karadibak, K., Gökmen, N., Erbayraktar, S., Göktay, Y., Taplu, A., Arkan, A. & Erkan, N. Effects of normovolaemic haemodilution on middle cerebral artery blood flow velocity and oxygen delivery, 330
Karalliedde, L. Cholinesterase estimations revisited: the clinical relevance, 313
Karamanlioğlu, B. *see* Memiş, D.
Karamanlioğlu, B. *see* Turan, A.
Karsli, B. *see* Eray, O.
Karsli, B. *see* Kayacan, N.
Katsikis, G. *see* Georgiou, M.
Kauranen, K. *see* Haavisto, E.
Kawachi, S. *see* Asai, T.
Kaya, F. N. *see* Yavaşçaoğlu, B.
Kaya, G. *see* Memiş, D.

Kayacan, N., Karsli, B., Zorlu, G., Döşemeci, L. & Erman, M. Hysteroscopic syndrome, 770
Kayarkar, R., Woolford, T. J. & Francis, G. A. Simple preoperative assessment to reduce the risk of traumatic epistaxis during nasotracheal intubation, 690
Kaymak, C., Gulhan, Y., Ozcan, A. O., Baltaci, B., Unal, N., Safak, M. A. & Oguz, H. Anaesthetic approach in a case of Goldenhar's syndrome, 836
Kelebek, N. *see* Yavaşcaoğlu, B.
Keller, C. *see* Loeckinger, A.
Kern, T. *see* Heid, F. M.
Khan, S. N. *see* Park, G. R.
Kiencke, P. *see* Kampe, S.
King, F. *see* Tucker, V.
Kinirons, B. *see* Gentili, M. E.
Kinsella, J. *see* Booth, M. G.
Klaghofer, R. *see* Hofer, C. K.
Klasen, J. *see* Junger, A.
Klein, A. A., Thoburn, C., Peters, C. & Hardee, P. S. G. F. Bronchial obstruction treated by fibreoptic-guided suction, 76
Kleinsasser, A. *see* Loeckinger, A.
Koçoğlu, H. *see* Pirbudak, L.
Kohjitani, A. & Obara, H. Subcutaneous epinephrine administration decreases lower oesophageal sphincter pressure and gastro-oesophageal pressure gradient in children under general anaesthesia, 189
Kong, S.-S. *see* Lin, T.-C.
Kontaratos, M. *see* Sator-Katzenschlager, S. M.
Koroglu, A. *see* Durmus, M.
Koroglu, A. *see* Togal, T.
Kose, Y. *see* Salihoglu, Z.
Kostopanagiotou, G. *see* Theodoraki, K.
Kotani, N., Taniuka, E., Tsubo, T., Ishihara, H. & Matsuki, A. Systemic heparinization during postoperative pulmonary embolism induces fatal complications, 382
Krafft, P. *see* Frietsch, T.
Krämer, B. K. *see* Ittner, K. P.
Krause, H. *see* Haeberle, H. A.
Kreuer, S. *see* Wilhelm, W.
Kubota, T. *see* Asai, T.
Kuhn, D. *see* Junger, A.
Kumle, B. *see* Piper, S. N.
Kurkal, P. *see* D'Mello, J.
Kurt, I. *see* Kurt, N.
Kurt, N., Kurt, I., Aygünes, B., Oral, H. & Tulunay, M. Effects of adding alfentanil or atracurium to lidocaine solution for intravenous regional anaesthesia, 522
Kuschinsky, W. *see* Frietsch, T.
Kutlay, O. *see* Yavaşcaoğlu, B.
Kyttä, J. V. *see* Tanskanen, P. E.

Laffey, J. G. *see* Sakamoto, H.
Lanham, P. *see* Brownlow, H.
Larsen, R. *see* Wilhelm, W.
Laxenaire, M.-C. *see* Mertes, P. M.
Le Corre, P. *see* Estebe, J. P.
Le Verge, R. *see* Estebe, J. P.
Lebuffe, G. & Vallet, B. A reply, 761
Leibundgut, D. L. *see* Sieber, T. J.
Lentschener, C. & Ozier, Y. Anaesthesia for elective liver resection: some points should be revisited, 780
Leroy, C. *see* Ricard-Hibon, A.
Leuwer, M. *see* Haeseler, G.
Levein, N.-G., Thörn, S.-E. & Wattwil, M. Effects of clonidine and dopamine on gastric tone, 99
Lin, T.-C., Hsu, C.-H., Kong, S.-S., Cherng, C.-H. & Wong, C.-S. Ventricular asystole and complete heart block after thoracic epidural analgesia for thymectomy, 460
Lindner, K. H. *see* Loeckinger, A.
Litscher, G. *see* Schwarz, G.
Lloyd, A. J. *see* Thompson, J. P.
Loeckinger, A., Hoermann, C., Keller, C., Lindner, K. H. & Kleinsasser, A. Augmented spontaneous breathing and pulmonary gas exchange during pneumoperitoneum, 424
Lönnqvist, P. A. *see* Naja, Z.
López, M. *see* Santiveri, X.
Loughrey, J. P. R., Walsh, F. & Gardiner, J. Prophylactic intravenous bolus ephedrine for elective Caesarean section under spinal anaesthesia, 63
Lowe, D. *see* O'Brien, B.
Lundberg, J. & Persson, K. A reply, 688

Mahla, E. *see* Gombotz, H.
Mahli, A., Coskun, D. & Akcali, D. T. Aetiology of convulsions due to stellate ganglion block: a review and report of two cases, 576
Maleck, W. H. *see* Piper, S. N.
Malhotra, S. K., Joshi, M., Grover, S., Sharma, S. C. & Dutta, A. Auditory function following spinal analgesia. Comparison of two spinal needles, 69
Mannaerts, G. H. H., Van Zundert, A. A. J., Meeusen, V. C. H., Martijn, H. & Rutten, H. J. T. Anaesthesia for advanced rectal cancer patients treated with combined major resections and intraoperative radiotherapy, 742
Marcus, M. A. E. *see* Brodner, G.
Margalit, N. *see* Friedman, Z.
Marmann, P. *see* Möllhoff, T.
Martijn, H. *see* Mannaerts, G. H. H.
Martínez, B. *see* Santiveri, X.
Marty, J. *see* Ricard-Hibon, A.
Matsukawa, T. *see* Noguchi, I.
Matsuki, A. *see* Hirota, K.
Matsuki, A. *see* Kotani, N.
Matsuki, A. *see* Tsubo, T.
Maue, D. *see* Haeseler, G.
Medved, I. *see* Šustić, A.
Meeus, P. *see* Tavan, R.
Meeusen, V. C. H. *see* Mannaerts, G. H. H.
Meignan, M. L. Morbidity in electroconvulsive therapy, 614
Mélot, C., Moraine, J. J. & Berré, J. Monitoring in neuroanaesthesia: update of clinical usefulness, 581
Memiş, D. *see* Turan, A.
Memiş, D., Turan, A., Karamanlioğlu, B., Kaya, G. & Pamukçu, Z. The prevention of propofol injection pain by tramadol or ondansetron, 47
Mertes, P. M. & Laxenaire, M.-C. Allergic reactions occurring during anaesthesia, 240
Metje, M. T. *see* Santiveri, X.
Metzler, H. *see* Gombotz, H.
Mielke, J. *see* Thornton, C.
Miller, D. M. & Pearce, A. C. A reply, 304
Miller, D. M. & Pearce, A. C. A reply, 306
Moerer, O., Baller, C., Hinz, J., Buscher, H. & Crozier, T. A. Neuromuscular effects of rapacuronium on the diaphragm and skeletal muscles in anaesthetized patients using cervical magnetic stimulation for stimulating the phrenic nerves, 883
Møller, A. M., Pedersen, T., Svendsen, P.-E. & Engquist, A. Perioperative risk factors in elective pneumonectomy: the impact of excess fluid balance, 57
Möllhoff, T., Schmidt, C., Van Aken, H., Berendes, E., Buerkle, H., Marmann, P., Reinbold, T., Prenger-Berninghoff, R., Tjan, T. D. T., Scheld, H. H. & Deng, M. C. Myocardial ischaemia in patients with impaired left ventricular function undergoing coronary artery bypass grafting – milrinone versus nifedipine, 796

Molyneux, M., Venn, R. & O'Dwyer, J. Localized abdominal pain following sympathetic blockade with bretylium for the management of complex regional pain syndrome, 147
Montpellier, D. *see* El Kettani, C.
Moore, E.W., Pollard, B. J. & Elliott, R. E. Anaesthetic agents in paediatric day case surgery: do they affect outcome?, 9
Moraine, J. J. *see* Mélot, C.
Morioka, T. *see* Fukui, K.
Mortier, E. *see* Cammu, G.
Mortier, E. P. *see* Struys, M. M. R. F.
Mostafa, S. M. & Murthy, B. V. S. Alcohol-associated admissions to an adult intensive care unit: an audit, 193
Murthy, B. V. S. *see* Mostafa, S. M.

Nagarathne, M. D. *see* Goonasekera, C. D. A.
Naja, Z., Ziade, M. F. & Lönnqvist, P. A. Bilateral paravertebral somatic nerve block for ventral hernia repair, 197
Nakayama, M., Yamamoto, J., Ichinose, H., Yamamoto, S., Kanaya, N. & Namiki, A. Effects of volume and concentration of lidocaine on epidural anaesthesia in pregnant females, 808
Namiki, A. *see* Nakayama, M.
Nathan, N. *see* Tecoult, E.
Necek, S. *see* Brock, H.
Nentwig, B. *see* Haeseler, G.
Neskovic, V. *see* Jankovic, Z.
Newton, D. E. F. *see* Thornton, C.
Ng, A., Habib, A., Swami, A., Smith, G., Nunns, D. & Davidson, A. C. Randomized controlled trial investigating the effect of transcervical papaverine and bupivacaine on postoperative analgesia following laparoscopic sterilization, 803
Niemitnen, P. *see* Vainionpää, V.
Nightingale, J. *see* Tucker, V.
Nishio, S. *see* Fukui, K.
Nobili, F. *see* Danelli, G.
Noguchi, I., Matsukawa, T., Ozaki, M. & Amemiya, Y. Propofol in low doses causes redistribution of body heat in male volunteers, 677
Nowicki, R. W. A. & Bull, P. T. Tongue piercing in an eclamptic patient, 844
Nunns, D. *see* Ng, A.

O'Brien, B., Lowe, D. & Awan, M. Anaesthetic implications of Aicardi's syndrome, 845
O'Donnell, B. & Synnott, A. Midodrine, an alternative to intravenous vasopressor therapy after spinal surgery, 841
O'Dwyer, J. *see* Molyneux, M.
Obara, H. *see* Kohjitani, A.
Oehmke, M. J. *see* Sator-Katzenschlager, S. M.
Oguz, H. *see* Kaymak, C.
Okawa, H. *see* Tsubo, T.
Oktay, C. *see* Eray, O.
Oral, H. *see* Kurt, N.
Ousmane, M. L., Fleyfel, M. & Vallet, B. Venous gas embolism during liver surgery with argon-enhanced coagulation, 225
Ozaki, M. *see* Noguchi, I.
Ozcan, A. O. *see* Kaymak, C.
Özcan, B. *see* Yavaşcaoğlu, B.
Ozgun, S. *see* Aydin, O. N.
Ozier, D. *see* Lentschener, C.

Pagedar, R. *see* D'Mello, J.
Pamuk, A. G. *see* Celebioğlu, B.
Pamukçu, Z. *see* Memiş, D.
Pamukçu, Z. *see* Turan, A.
Pandey, K. *see* D'Mello, J.
Papadimitriou, J. *see* Theodoraki, K.
Papadopoulos, G., Sgouropoulou, S., Arnaoutoglou, E. & Petrou, A. Postoperative hypoxaemia in a patient with patent foramen ovale, 152
Papagiannopoulou, P. *see* Georgiou, M.
Papaziogas, B. *see* Vretzakis, G.
Pappachan, J. *see* Brownlow, H. A.
Park, G. R. & Khan, S. N. Murder and the ICU, 621
Parlapani, A. *see* Georgiou, M.
Parpaglioni, R., Capogna, G., Celleno, D. & Fusco, P. Intraoperative fetal oxygen saturation during Caesarean section: general anaesthesia using sevoflurane with either 100% oxygen or 50% nitrous oxide in oxygen, 115
Paşaoğlu, I. *see* Celebioğlu, B.
Pasvol, G. *see* Thornton, C.
Pearce, A. C. *see* Miller, D. M.
Pederneiras, S. G. *see* de Oliveira Filho, G. R.
Pedersen, T. *see* Møller, A. M.
Pedicini, M. S. *see* Iannuzzi, E.
Pennefather, S. H. *see* Guha, A.
Perel, A. *see* Friedman, Z.
Persson, K. *see* Lundberg, J.
Peters, C. *see* Klein, A. A.
Pethiyagoda, C. J. B. *see* Goonasekera, C. D. A.
Petrou, A. *see* Papadopoulos, G.
Piepenbrock, S. *see* Haeseler, G.
Piepgras, A. *see* Frietsch, T.
Pietropaoli, P. *see* Della Rocca, G.
Piper, S. N., Suttner, S. W., Maleck, W. H., Kumle, B., Haisch, G. & Boldt, J. Effects of sodium nitroprusside-induced controlled hypotension on pancreatic function assessed by pancreatitis-associated protein in patients undergoing radical prostatectomy, 609
Pirbudak, L., Tuncer, S., Koçoğlu, H., Göksu, S. & Çelik, Ç. Fentanyl added to bupivacaine 0.05% or ropivacaine 0.05% in patient-controlled epidural analgesia in labour, 271
Plaja, I. *see* Santiveri, X.
Pollard, B. J. *see* Moore, E. W.
Pompei, L. *see* Della Rocca, G.
Popat, M. *see* Caponas, G.
Prachalias, A. *see* Theodoraki, K.
Prenger-Berninghoff, R. *see* Möllhoff, T.
Pyrsopoulos, N. *see* Theodoraki, K.

Radojevic, D. *see* Jankovic, Z.
Randell, T. T. *see* Tanskanen, P. E.
Raucoules-Aime, M. *see* Dubost, T.
Reckziegel, G., Friederich, P. & Urban, B. W. Ketamine effects on human neuronal Na^+ channels, 634
Redmond, H. P. *see* Tyther, R.
Rehak, P. *see* Gombotz, H.
Rehberg, B. *see* Bruhn, J.
Reinbold, T. *see* Möllhoff, T.
Ricard-Hibon, A., Chollet, C., Leroy, C. & Marty, J. Succinylcholine improves the time of performance of a tracheal intubation in prehospital critical care medicine, 361
Riess, M. L., Graefe, U. A., Goeters, C., Van Aken, H. & Bone, H. G. Sedation assessment in critically ill patients with bispectral index, 18
Robays, H. M. *see* Struys, M. M. R. F.
Rogers, S. A. *see* Guha, A.
Rohlfs, R. *see* Filipovic, M.
Röhrig, R. *see* Junger, A.
Rolfseng, O. K., Skogvoll, E. & Borchgrevink, P. C. Epidural bupivacaine with sufentanil or fentanyl during labour: a randomized, double-blind study, 812
Ronco, P. M. *see* Cittanova, M. L.
Röpcke, H. *see* Bruhn, J.
Rossaint, R. *see* Schulz-Stübner, S.
Rosseel, P. M. J. *see* El Azab, S. R.
Rothe, G. *see* Fröhlich, D.

Rowbotham, D. J. *see* Sharpe, P.
Rushe, C. *see* Halpenny, M.
Rushman, S. C. *see* Thompson, J. P.
Russell, G. N. *see* Guha, A.
Rutten, H. J. T. *see* Mannaerts, G. H. H.

Saatweber, P. *see* Adams, H. A.
Sacerdoti, G. *see* Iannuzzi, E.
Safak, M. A. *see* Kaymak, C.
Şahin, N. *see* Kara, H.
Sakai, I. *see* Tsubo, T.
Sakamoto, H., Corcoran, T. B., Laffey, J. G. & Shorten, G. D. Isoprostanes – markers of ischaemia reperfusion injury, 550
Salihoglu, Z., Demiroluk, S., Demirkiran, O. & Kose, Y. Comparison of effects of remifentanil, alfentanil and fentanyl on cardiovascular responses to tracheal intubation in morbidly obese patients, 125
Sanders, M. K. & Wee, M. Y. K. Multicompartment block in a parturient undergoing urgent Caesarean section, 226
Santamaria, L. B. *see* Fodale, V.
Santiveri, X., Arxer, A., Plaja, I., Metje, M. T., Martínez, B., Villalonga, A. & López, M. Anaesthetic and postoperative analgesic effects of spinal clonidine as an additive to prilocaine in the transurethral resection of urinary bladder tumours, 589
Sato, N. *see* Takeda, S.
Sator-Katzenschlager, S. M., Oehmke, M. J., Kontaratos, M., Wedrich, A., Heinze, G. & Weinstabl, C. Effect of different doses of cisatracurium on intraocular pressure in sedated patients, 823
Scawn, N. D. A. *see* Guha, A.
Scheffer, G. J. *see* El Azab, S. R.
Scheld, H. H. *see* Möllhoff, T.
Schmidt, C. *see* Möllhoff, T.
Schmidt, R. *see* Haeberle, H. A.
Schmitz, C. S. *see* Adams, H. A.
Schmitz, G. *see* Fröhlich, D.
Schmitz, T. P. *see* Friederich, P.
Scholz, M. *see* von Spiegel, T.
Schorn, B. *see* von Spiegel, T.
Schraag, S. & Flaschar, J. Delivery performance of commercial target-controlled infusion devices with Diprifusor® module, 357
Schulte am Esch, J. *see* Freitag, M.
Schulte am Esch, J. *see* Gerbershagen, M. U.
Schultz, U. & Buhre, W. Investigating the usefulness of the laryngeal tube in spontaneously breathing patients, 303
Schulz-Stübner, S., Zielske, D. & Rossaint, R. Comparison between nasal and intravenous desmopressin for the treatment of aminosalicylic acid-induced platelet dysfunction, 647
Schummer, C. *see* Schummer, W.
Schummer, W. & Schummer, C. Another neurological complication of central venous cannulation, 73
Schummer, W. & Schummer, C. Central venous catheter malposition in the azygos arch, 832
Schurig, V. *see* Haeberle, H. A.
Schwab, S. *see* Frietsch, T.
Schwarz, G. & Litscher, G. Transcranial cerebral oximetry, transcranial Doppler sonography, and heart rate variability: useful neuromonitoring tools in anaesthesia and intensive care?, 543
Schwilden, H. *see* Ihmsen, H.
Schywalsky, M. *see* Ihmsen, H.
Seeberger, M. D. *see* Filipovic, M.
Segal, E. *see* Friedman, Z.
Sgouropoulou, S. *see* Papadopoulos, G.
Sharma, S. C. *see* Malhotra, S. K.
Sharpe, P., Asif, M., Victoria, A. & Rowbotham, D. J. Iontophoretically applied lidocaine and the prevention of pain associated with the injection of intravenous propofol: a comparison with intravenous lidocaine, 170
Shimoji, K. *see* Taga, K.
Shipton, E. A. *see* Wisely, N. A.
Shorten, G. *see* Tyther, R.
Shorten, G. D. *see* Halpenny, M.
Shorten, G. D. *see* Sakamoto, H.
Sieber, T. J. & Leibundgut, D. L. Operating room management and strategies in Switzerland: results of a survey, 415
Šimić, O. *see* Šustić, A.
Simpson, M. E. *see* Conway, D. H.
Skarvan, K. *see* Filipovic, M.
Skogvoll, E. *see* Rolfseng, O. K.
Slappendel, R., Weber, E. W. G., Benraad, B., Dirksen, R. & Bugter, M. L. T. Does ibuprofen increase perioperative blood loss during hip arthroplasty?, 829
Smith, G. *see* Ng, A.
Smyrniotis, V. *see* Theodoraki, K.
Sokolovic, E., Biro, P., Wyss, P., Werthemann, C., Haller, U., Spahn, D. & Szucs, T. Impact of the reduction of anaesthesia turnover time on operating room efficiency, 560
Somers, A. A. L. *see* Struys, M. M. R. F.
Spahn, D. *see* Sokolovic, E.
Stamenkovic, D. *see* Jankovic, Z.
Standl, T. *see* Freitag, M.
Struys, M. *see* Cammu, G.
Struys, M. M. R. F., Somers, A. A. L., Van Den Eynde, N., Herregods, L. L. G., Dyzers, D., Robays, H. M. & Mortier, E. P. Cost-reduction analysis of propofol versus sevoflurane: maintenance of anaesthesia for gynaecological surgery using the bispectral index, 727
Studer, W. *see* Filipovic, M.
Šustić, A., Medved, I. & Šimić, O. Ultrasound-guided placement of intra-aortic balloon pump, 149
Sutherland, P. *see* Brownlow, H.
Suttner, S. W. *see* Piper, S. N.
Suzuki, A. *see* Tsubo, T.
Svendsen, P.-E. *see* Møller, A. M.
Swami, A. *see* Ng, A.
Synnott, A. *see* O'Donnell, B.
Szucs, T. *see* Sokolovic, E.

Taeger, K. *see* Fröhlich, D.
Taeger, K. *see* Ittner, K. P.
Taga, K., Fujihara, H., Baba, H., Yamakura, T. & Shimoji, K. Workforce and regional distribution of anaesthesiologists in Japan, 530
Tagariello, V. *see* De Tommaso, O.
Takahashi, S. *see* Fukui, K.
Takeda, S. & Tomaru, T. Haemodynamic and splanchnic organ blood-flow responses to hypotension induced by the calcitonin gene-related, 876
Takeda, S., Sato, N. & Tomaru, T. Haemodynamic and splanchnic organ blood flow responses during sevoflurane-induced hypotension in dogs, 442
Takenaka-Hamaya, C., Hamaya, Y. & Dohi, S. Epidural morphine injection after combined spinal and epidural anaesthesia, 672
Tanioka, F. *see* Kotani, N.
Tanskanen, P. E., Kyttä, J. V. & Randell, T. T. QT interval and QT dispersion during the induction of anaesthesia in patients with subarachnoid haemorrhage: a comparison of thiopental and propofol, 749
Taplu, A. *see* Karadibak, K.
Tavan, R., Coronel, B., Duperret, S., Meeus, P. & Blondet, R. Peroperative opening of a patent foramen ovale caused by

pulmonary artery obstruction due to knotted pulmonary artery catheter, 145
Tchaoussoff, J. *see* El Kettani, C.
Tecoult, E. & Nathan, N. A reply, 615
Theodoraki, K., Kostopanagiotou, G., Smyrniotis, V., Arkadopoulos, N., Prachalias, A., Pyrsopoulos, N. & Papadimitriou, J. Haemodynamic changes in ischaemic vs. anhepatic pig experimental model of acute liver failure, 40
Thoburn, C. *see* Klein, A. A.
Thompson, J. P., Rushman, S. C., Fox, A. J., Lloyd, A. J. & Atcheson, R. A. Using the Sustained Attention to Response Test to detect cognitive dysfunction after day case surgery, 585
Thörn, S.-E. *see* Levein, N.-G.
Thorniley, M. *see* Thornton, C.
Thornton, C., Heyderman, R. S., Thorniley, M., Curtis, N., Mielke, J., Pasvol, G. & Newton, D. E. F. Auditory- and somatosensory-evoked potentials in cerebral malaria and anaesthesia: a comparison, 717
Tjan, T. D. T. *see* Möllhoff, T.
Togal, T. *see* Durmus, M.
Togal, T., Durmus, M., Koroglu, A., Demirbilek, S., Karaaslan, K. & Ersoy, O. Anaesthesia for Caesarean section in the presence of aortic coarctation, 768
Tokarz, A., Gaszyński, T., Gaszyński, W. & Arkuszewski, P. General anaesthesia for a child with Rubinstein-Taybi syndrome, 896
Tokarz, A., Gaszyński, T., Gaszyński, W. & Arkuszewski, P. General anaesthesia with remifentanil and propofol for a patient with centronuclear (myotubular) myopathy, 842
Tomaru, T. *see* Takeda, S.
Toprak, H. I. *see* Durmus, M.
Torri, G. *see* Casati, A.
Torri, G. *see* Danelli, G.
Troisi, R. *see* Cammu, G.
Tsubo, T. *see* Kotani, N.
Tsubo, T., Sakai, I., Suzuki, A., Okawa, H., Ishihara, H. & Matsuki, A. Lung image changes during bronchopulmonary lavage estimated using transoesophageal echocardiography, 141
Tucci, M. *see* Hofer, C. K.
Tucker, V., King, F. & Nightingale, J. Thoracic epidurals and avoidance of potential neurological damage, 772
Tulunay, M. *see* Kurt, N.
Tuncer, S. *see* Pirbudak, L.
Turan, A. *see* Memiş, D.
Turan, A., Memiş, D., Karamanlioğlu, B., Çolak, A., Pamukçu, Z. & Turan, N. Effect of aminophylline on recovery from sevoflurane anaesthesia, 452
Turan, N. *see* Turan, A.
Turkoz, A. *see* Durmus, M.
Tyagi, A. & Bhattacharya, A. Central neuraxial blocks and anticoagulation: a review of current trends, 317
Tylman, M., Arnestad, J.-P., Bengtsson, A. & Bengtson, J.-P. Salvage with a haemofiltration technique unsuitable in hip prosthesis surgery due to free haemoglobin, 766
Tyther, R., Halligan, M., Wang, J., Redmond, H. P. & Shorten, G. Effects of chronic occupational exposure to anaesthetic gases on the rate of neutrophil apoptosis among anaesthetists, 604
Tzabazis, A. *see* Ihmsen, H.

Uçkunkaya, N. *see* Yavaşcaoğlu, B.
Ulusan, V. *see* Kara, H.
Unal, N. *see* Kaymak, C.
Ünlüsoy, E. Ö. *see* Karaca, S.
Urban, B. W. *see* Reckziegel, G.
Uysalel, A. *see* Aşık, İ.

Vainionpää, V., Haavisto, E. & Nieminen, P. Minor haemodynamic responses in coronary bypass patients following tracheal intubation using either a lightwand device (Trachlight®) or direct laryngoscopy, 229
Vallet, B. *see* Lebuffe, G.
Vallet, B. *see* Ousmane, M. L.
Van, D. M. *see* Gozal, Y.
Van Aken, H. *see* Brodner, G.
Van Aken, H. *see* Möllhoff, T.
Van Aken, H. *see* Riess, M. L.
Van Den Eynde, N. *see* Struys, M. M. R. F.
van Gessel, E. *see* Hofer, C. K.
Van Gessel, E. *see* Weber, A.
van Strik, R. *see* El Azab, S. R.
van Wijk, E. M. *see* El Azab, S. R.
Van Winkle, D. M. *see* Gozal, Y.
Van Zundert, A. A. J. *see* Mannaerts, G. H. H.
Venn, R. *see* Molyneux, M.
Vermeyen, K. M. *see* Himpe, D. G.
Verpont, M.-C. *see* Cittanova, M. L.
Verton, C. *see* Gentili, M. E.
Vicenzi, M. *see* Gombotz, H.
Victoria, A. *see* Sharpe, P.
Villalonga, A. *see* Santiveri, X.
Vollmer, T., Genzwuerker, H. V. & Ellinger, K. Fibreoptic control of the laryngeal tube position, 306
von Breunig, F. *see* Gerbershagen, M. U.
von Spiegel, T., Giannaris, S., Schorn, B., Scholz, M., Wietasch, G. J. K. & Hoeft, A. Effects of induction of anaesthesia with sufentanil and positive-pressure ventilation on the intra- to extrathoracic volume distribution, 428
Vretzakis, G., Ferdi, E. & Papaziogas, B. Insidious hypothyroidism unmasked after operation, 532

Wahbe, F. *see* Cittanova, M. L.
Wahl, H. G. *see* Haeberle, H. A.
Walsh, F. *see* Loughrey, J. P. R.
Wang, J. *see* Tyther, R.
Wappler, F. *see* Gerbershagen, M. U.
Waschke, K. F. *see* Frietsch, T.
Watson, S., Calam, R. & Jimmieson, P. Can computers help in assessing children's postoperative pain? Initial validation of a computer-assisted interview, 510
Wattwil, M. *see* Levein, N.-G.
Weber, A., Fournier, R., Van Gessel, E. & Gamulin, Z. Sciatic nerve block and the improvement of femoral nerve block analgesia after total knee replacement, 834
Weber, E. W. G. *see* Slappendel, R.
Wedrich, A. *see* Sator-Katzenschlager, S. M.
Wee, M. Y. K. *see* Sanders, M. K.
Weinstabl, C. *see* Sator-Katzenschlager, S. M.
Weißhorn, R. *see* Gerbershagen, M. U.
Werthemann, C. *see* Sokolovic, E.
Westphal, M. *see* Brodner, G.
Wietasch, G. J. K. *see* von Spiegel, T.
Wild, B., Eschweiler, G. W. & Bartels, M. Morbidity in electroconvulsive therapy, 616
Wildsmith, J. A. W. Oasis or mirage? The safety of outpatient dental anaesthesia in hospital, 762
Wilhelm, S. *see* Freitag, M.
Wilhelm, W., Biedler, A., Huppert, A., Kreuer, S., Bücheler, O., Ziegenfuss, T. & Larsen, R. Comparison of the effects of remifentanil or fentanyl on anaesthetic induction characteristics of propofol, thiopental or etomidate, 350
Wisely, N. A. & Shipton, E. A. Long QT syndrome and anaesthesia, 853
Wittmann, S. *see* Fröhlich, D.
Wolff, R. A. *see* Gozal, Y.
Wolkiewiez, J. *see* Dubost, T.

Wong, C.-S. *see* Lin, T.-C.
Woolford, T. J. *see* Kayarkar, R.
Wyss, P. *see* Sokolovic, E.

Yamakura, T. *see* Taga, K.
Yamamoto, J. *see* Nakayama, M.
Yamamoto, S. *see* Nakayama, M.
Yavaşcaoğlu, B., Acar, Ş. H. V., Kahveci, F., Kaya, F. N. & Özcan, B. Cuffed oropharyngeal airway as a suitable alternative to the laryngeal mask airway for minor outpatient surgery, 203
Yavaşcaoğlu, B., Cebelli, V., Kelebek, N., Uçkunkaya, N. & Kutlay, O. Comparison of different priming techniques on the onset time and intubating conditions of rocuronium, 517
Yener, F. *see* Celebioğlu, B.

Ziade, M. F. *see* Naja, Z.
Ziegenfuss, T. *see* Wilhelm, W.
Zielske, D. *see* Schulz-Stübner, S.
Zimmermann, M. *see* Ittner, K. P.
Zollinger, A. *see* Hofer, C. K.
Zorlu, G. *see* Kayacan, N.